

Case Number:	CM14-0187968		
Date Assigned:	11/18/2014	Date of Injury:	12/08/2012
Decision Date:	10/30/2015	UR Denial Date:	11/05/2014
Priority:	Standard	Application Received:	11/11/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This is a 47 year old male, whose date of injury was December 8, 2012. The medical records indicated the injured worker was treated for left knee pain, neck pain and low back pain. According to the progress notes on 10-14-2014 the injured worker reported knee pain, which he rated a 9 on a 10-point scale. His medications included Aleve four per day. His previous medications included Lidocaine, Ultram and Norco, which helped. On physical examination, he had diminished range of motion of the cervical spine. Left lateral rotation and extension increased his pain with some numbness and tingling down the left arm. He had full range of motion of the shoulders with no tenderness to palpation. He had diminished range of motion of the lumbar spine particularly with flexion and extension. He had tenderness to palpation on the left low back and buttock area. Straight leg raise was positive on the left. His left knee had swelling and he was not able to fully extend the knee. Flexion was only at 80 degrees and ligament stress test increased his symptoms. The injured worker was diagnosed as having chronic neck and left upper extremity pain, chronic low back pain, and chronic left knee pain. A request for authorization for Ultracet 37.5 - 325 mg of unspecified quantity 2-4 per day and for Lidoderm patches for the left knee was received on October 29, 2014. On November 5, 2014, the Utilization Review physician determined Ultracet 37.5 - 325 mg of unspecified quantity 2-4 per day and for Lidoderm patches for the left knee was not medically necessary based on the CA MTUS Chronic Pain Medical Treatment Guidelines.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Ultracet 37.5/325 mg of unspecified quantity (Rx 10/14/14): Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids for chronic pain.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain= patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the 4 A's (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids: (a) If the patient has returned to work, (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function or how the medication improves activities. The work status is not mentioned. There is also no quantity specified in the request. Therefore all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.

Lidoderm patches of unspecified quantity (R 10/14/14): Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: The California chronic pain medical treatment guidelines section on topical lidocaine states: Lidocaine Indication: Neuropathic pain Recommended for localized peripheral pain after there has been evidence of a trial of first-line therapy (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). Topical lidocaine, in the formulation of a dermal patch (Lidoderm) has been designated for orphan status by the FDA for neuropathic pain. Lidoderm is also used off-label for diabetic neuropathy. No other commercially approved topical formulations of lidocaine (whether creams, lotions or gels) are indicated for neuropathic pain. Non-dermal patch formulations are generally indicated as local anesthetics and anti-pruritics. Further research is needed to recommend this treatment for chronic neuropathic pain disorders other than post-herpetic neuralgia. Formulations that do not involve a dermal-patch system are generally indicated as local anesthetics and anti-pruritics. In February 2007 the FDA notified consumers and healthcare professionals of the potential hazards of the use of topical lidocaine. Those at particular risk were individuals that applied large amounts of this substance over large areas, left the products on for long periods of time, or used the agent with occlusive dressings. Systemic exposure was highly variable among patients. Only FDA-approved products are currently recommended. (Argoff, 2006) (Dworkin, 2007) (Khaliq-Cochrane, 2007) (Knotkova, 2007) (Lexi-Comp, 2008) Non-neuropathic pain: Not recommended. There is only one trial that tested 4% lidocaine for treatment of chronic muscle pain. The results showed there was no superiority over placebo. (Scudds, 1995) This medication is recommended for localized peripheral pain. The patient has lumbago. There is no documentation of failure of first line neuropathic pain medications. Therefore, criteria as set forth by the California MTUS as outlined above have not been met and the request is not medically necessary.