MAXIMUS Federal Services

Case Number:	CM15-0126289		
Date Assigned:	07/16/2015	Date of Injury:	08/28/1990
Decision Date:	10/08/2015	UR Denial Date:	06/15/2015
Priority:	Standard	Application Received:	06/30/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: California
Certification(s)/Specialty: Emergency Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 58 year old male, who sustained an industrial injury on 8/28/90. The injured worker has complaints of low back pain and neck pain. The documentation noted that he injured worker had restricted mobility at the left shoulder where he tolerated only about 90 degrees of active abduction and there was trigger point myofascial tenderness around the cervical paraspinous muscles and tenderness around the left shoulder joint. The diagnoses have included spinal stenosis in cervical region; cervical spondylosis; lumbar spondylosis and chronic pain syndrome. Treatment to date has included knee surgery times two; acupuncture; cervical pillow; home exercise program and physical therapy. The request was for flubriprofen powder 6gm; lidocaine 1.5gm; varasapro base cream 22.5mg; gabapentin powder 3gm; amitriptyline 1.5gm; capsaicin 0.0075gm; varsapro base cream 25.49gm; cyclobenzaprine powder 3 gm; lidocaine 0.6gm; diclofenac 100mg #60 and omeprazole 20mg #60.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Flurbiprofen Powder 6gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: The request is for the use of a topical NSAID for pain relief. There are specific criteria require for use based on the guidelines. The MTUS states the following: The efficacy in clinical trials for this treatment modality has been inconsistent and most studies are small and of short duration. Topical NSAIDs have been shown in meta-analysis to be superior to placebo during the first 2 weeks of treatment for osteoarthritis, but either not afterward, or with a diminishing effect over another 2-week period. (Lin, 2004) (Bjordal, 2007) (Mason, 2004) When investigated specifically for osteoarthritis of the knee, topical NSAIDs have been shown to be superior to placebo for 4 to 12 weeks. Indications: Osteoarthritis and tendinitis, in particular, that of the knee and elbow or other joints that are amenable to topical treatment: Recommended for short-term use (4-12 weeks). There is little evidence to utilize topical NSAIDs for treatment of osteoarthritis of the spine, hip or shoulder. FDA-approved agents: Voltaren Gel 1% (diclofenac): Indicated for relief of osteoarthritis pain in joints that lend themselves to topical treatment (ankle, elbow, foot, hand, knee, and wrist). It has not been evaluated for treatment of the spine, hip or shoulder. In this case, as indicated above, the patient would not qualify for the use of this medication based on the treatment duration. As such, the request is not medically necessary.

Lidocaine 1.5gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: The request is for the use of topical lidocaine. The MTUS guidelines state the following: Lidocaine Indication: Neuropathic pain: Recommended for localized peripheral pain after there has been evidence of a trial of first-line therapy (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). Topical lidocaine, in the formulation of a dermal patch (Lidoderm) has been designated for orphan status by the FDA for neuropathic pain. Lidoderm is also used off-label for diabetic neuropathy. No other commercially approved topical formulations of lidocaine (whether creams, lotions or gels) are indicated for neuropathic pain. Non-dermal patch formulations are generally indicated as local anesthetics and anti-pruritics. Further research is needed to recommend this treatment for chronic neuropathic pain disorders other than post-herpetic neuralgia. In this case, as stated above, the patient does not meet the criteria for use of this product in this formulation. There is a requirement of documentation of a first-line therapy trial prior to use of a lidocaine dermal patch. There is also no other commercially approved topical formulations of lidocaine indicated for neuropathic pain other than Lidoderm. As such, the request is not medically necessary.

Varsapro Base Cream 22.5gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: The request is for the use of a compounded medication for topical use to aid in pain relief. These products contain multiple ingredients which each have specific properties and mechanisms of action. The MTUS guidelines state the following: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. In this case, the use of this product is not indicated. This is due to poor scientific evidence regarding efficacy for the patient's condition. As such, the request is not medically necessary.

Gabapentin Powder 3gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: The request is for the use of a compounded medication for topical use to aid in pain relief. These products contain multiple ingredients which each have specific properties and mechanisms of action. The MTUS guidelines state the following: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. In this case, the use of gabapentin is stated to be not indicated for use for the patient's condition. The guidelines state the following: Gabapentin: Not recommended. There is no peer-reviewed literature to support use. As such, the request is not medically necessary.

Amitriptyline 1.5gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: The request is for the use of a compounded medication for topical use to aid in pain relief. These products contain multiple ingredients which each have specific properties and mechanisms of action. The MTUS guidelines state the following: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. In this case, the use of this product is not indicated. This is secondary to poor scientific evidence of efficacy for the patient's condition when applied topically. As such, the request is not medically necessary.

Capsaicin 0.0075gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: The request is for the use of a compounded medication for topical use to aid in pain relief. These products contain multiple ingredients which each have specific properties and mechanisms of action. The MTUS guidelines state the following: "Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended." In this case, the compounded topical treatment contains Capsaicin. Qualifying factors for this product is indicated by the following per the guidelines: Capsaicin: Recommended only as an option in patients who have not responded or are intolerant to other treatments. Formulations: Capsaicin is generally available as a 0.025% formulation (as a treatment for osteoarthritis) and a 0.075% formulation (primarily studied for post-herpetic neuralgia, diabetic neuropathy and post-mastectomy pain). There have been no studies of a 0.0375% formulation of capsaicin and there is no current indication that this increase over a 0.025% formulation would provide any further efficacy. Indications: There are positive randomized studies with capsaicin cream in patients with osteoarthritis, fibromyalgia, and chronic non-specific back pain, but it should be considered experimental in very high doses. Although topical capsaicin has moderate to poor efficacy, it may be particularly useful (alone or in conjunction with other modalities) in patients whose pain has not been controlled successfully with conventional therapy. The number needed to treat in musculoskeletal conditions was 8.1. The number needed to treat for neuropathic conditions was 5.7. (Robbins, 2000) (Keitel, 2001) (Mason-BMJ, 2004) In this case, as stated above, the patient would not qualify for the use of capsaicin based on the diagnosis. As such, the request is not medically necessary.

Varsapro Base Cream 25.49gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: The request is for the use of a compounded medication for topical use to aid in pain relief. These products contain multiple ingredients which each have specific properties and mechanisms of action. The MTUS guidelines state the following: "Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended." In this case, the use of this product is not indicated. This is due to poor scientific evidence regarding efficacy for the patient's condition. As such, the request is not medically necessary.

Cyclobenzaprine Powder 3gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: The request is for the use of a compounded medication for topical use to aid in pain relief. These products contain multiple ingredients which each have specific properties and mechanisms of action. The MTUS guidelines state the following: "Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended." In this case, the use of the topical muscle relaxant is not indicated for use for the patient's condition. The MTUS states the following: "There is no evidence for use of any other muscle relaxant as a topical product." As such, the request is not medically necessary.

Lidocaine 0.6gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: The request is for the use of topical lidocaine. The MTUS guidelines state the following: Lidocaine Indication: Neuropathic pain: Recommended for localized peripheral pain after there has been evidence of a trial of first-line therapy (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). Topical lidocaine, in the formulation of a dermal patch (Lidoderm) has been designated for orphan status by the FDA for neuropathic pain. Lidoderm is also used off-label for diabetic neuropathy. No other commercially approved topical formulations of lidocaine (whether creams, lotions or gels) are indicated for neuropathic pain. Non-dermal patch formulations are generally indicated as local anesthetics and anti-pruritics. Further research is needed to recommend this treatment for chronic neuropathic pain disorders other than post-herpetic neuralgia. In this case, as stated above, the patient does not meet the criteria for use of this product in this formulation. There is a requirement of documentation of a first-line therapy trial prior to use of a lidocaine dermal patch. There is also no other commercially approved topical formulations of lidocaine indicated for neuropathic pain other than Lidoderm. As such, the request is not medically necessary.

Diclofenac 100mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG); Pain Chapter.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic)/NSAIDs (non-steroidal anti-inflammatory drugs).

Decision rationale: The request is for the use of a medication in the NSAID class. The ODG state the following regarding this topic: Specific recommendations: Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008) Back Pain-Acute low back pain & acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting to negative evidence that NSAIDs are more effective than acetaminophen for acute LBP. (van Tulder, 2006) (Hancock, 2007) For patients with acute low back pain with sciatica a recent Cochrane review (including three heterogeneous randomized controlled trials) found no differences in treatment with NSAIDs vs. placebo. In patients with axial low back pain this same review found that NSAIDs were not more effective than acetaminophen for acute low-back pain, and that acetaminophen had fewer side effects. (Roelofs-Cochrane, 2008) The addition of NSAIDs or spinal manipulative therapy does not appear to increase recovery in patients with acute low back pain over that received with acetaminophen treatment and advice from their physician. (Hancock, 2007) Back Pain-Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another. (Roelofs-Cochrane, 2008) See also Anti-inflammatory medications. Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long-term neuropathic pain, but they may be useful to treat breakthrough pain and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in patients with neuropathic pain. (Namaka, 2004) (Gore, 2006) See NSAIDs, GI symptoms & cardiovascular risk; NSAIDs, hypertension and renal function; & Medications for acute pain (analgesics). Besides the above well-documented side effects of NSAIDs, there are other less well-known effects of NSAIDs, and the use of NSAIDs has been shown to possibly delay and hamper healing in all the soft tissues, including muscles, ligaments, tendons, and cartilage. (Maroon, 2006) The risks of NSAIDs in older patients, which include increased cardiovascular risk and gastrointestinal toxicity, may outweigh the benefits of these medications. (AGS, 2009) As stated above, acetaminophen would be considered first-line treatment for chronic pain. In this case, the use of an NSAID is not advised. This is secondary

to the duration of use and significant side effect profile. Also, the use of NSAIDs is known to delay the healing of soft tissue including ligaments, tendons, and cartilage. As such, the request is not medically necessary.

Omeprazole 20mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: The request is for the use of a medication in the class of a proton pump inhibitor. It is indicated for patients with peptic ulcer disease. It can also be used as a preventative measure in patients taking non-steroidal anti-inflammatories for chronic pain. Unfortunately, they do have certain side effects including gastrointestinal disease. The MTUS guidelines states that patients who are classified as intermediate or high risk, should be treated prophylactically. Criteria for risk are as follows: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Due to the fact the patient does not meet to above stated criteria, the request for use is not medically necessary.